

This exam has 5 problems. Please start each new problem at the top of a new page in your blue book. Show all your work in your blue book and simplify your answers. Answers with no justification will receive no points. You are allowed one 8.5×11-in page of notes (ONE side). NO calculators, smartphones/watches, or any other electronic devices allowed.

Problem 1 (17 pts)

A wire lies along the intersection of the surfaces  $z = \frac{1}{3}xy$  and  $y = \frac{1}{2}x^2$  from  $(0;0;0)$  to  $(4;8;\frac{32}{3})$ . Suppose the charge density at any point on the wire is given by

$$(x; y; z) = x \frac{\text{Coulombs}}{\text{meter}}$$

Find the total charge on the wire. (You can assume distance in  $xyz$ -space is measured in meters).

SOLUTION:

$$\text{Charge} = \int_C (x; y; z) ds = \int_C x ds$$

where  $C$  is the path traced out by the wire.

To evaluate, we first need to parameterize the wire.


One simple parameterization:

$$\mathbf{r}(t) = \langle ht; \frac{1}{2}t^2; \frac{1}{6}t^3 \rangle; \quad 0 \leq t \leq 4$$

$$\text{Charge} = \int_0^4 \|\mathbf{r}'(t)\| dt$$

$$\mathbf{r}'(t) = \langle h; t; \frac{1}{2}t^2 \rangle$$


=



(b) We will use the following substitution:

$$u = \frac{x}{y}$$

$$v = x + y$$



Solving for the  $xy$ -transformation yields:

$$x = \frac{uv}{1+u}$$

$$y = \frac{v}{1+u}$$

NextmQ [(Ne)omput250(the)-250(foJJacobia)TJ/F53 10.9091 Tf 158.5759620.6052d [(v)]@J/F30 10.9091 Tf 8.768


**XV**

Let  $g(x; y; z) = xy - z$

Since the surface is given as a function of  $x$  and  $y$  it's easiest to project the surface onto the  $xy$ -plane. Thus, let  $\hat{p} = \mathbf{k}$ .

$$dS = \frac{|\mathbf{r}_x \cdot \mathbf{g}_x + \mathbf{r}_y \cdot \mathbf{g}_y|}{|\mathbf{j}|} dA = \frac{\sqrt{y^2 + x^2 + 1}}{1} dA$$

$$T_{ave} = \frac{\iint_S 3xyz \, dS}{\iint_S dS} = \frac{\iint_R 3xyz \sqrt{x^2 + y^2 + 1} \, dA}{\iint_R \sqrt{x^2 + y^2 + 1} \, dA}$$



(b)



(c)

(d) (i)

$$\text{Mass} = \int_0^a \int_0^{\sqrt{a^2 - z^2}} \int_0^{\sqrt{a^2 - z^2 - y^2}} \rho \sqrt{x^2 + y^2} \, dz \, dx \, dy + \int_0^a \int_0^y \int_0^a \rho \sqrt{x^2 + y^2} \, dz \, dx \, dy$$

(ii)

$$\text{Mass} = \int_0^a \int_0^{\sqrt{a^2 - z^2}} \int_0^{\sqrt{a^2 - z^2 - y^2}} \rho \sqrt{x^2 + y^2} \, dz \, dx \, dy + \int_0^a \int_0^y \int_0^a \rho \sqrt{x^2 + y^2} \, dz \, dx \, dy$$

Problem 5 (17 points)

A neighborhood sits in the region in the  $xy$ -plane given by

$$x^2 + y^2 \leq 4 \text{ and } x \geq 1$$

(where distances in the  $xy$ -plane are measured in miles). An earthquake occurs with epicenter at the origin. Suppose at each point in the neighborhood, the energy density released from the earthquake is given by the function

$$E(x; y) = \frac{10^6}{(d(x; y))^3} \frac{\text{joules}}{\text{miles}^2}$$

where  $d(x; y)$  is the distance from  $(x; y)$  to the epicenter.

Find the total amount of energy (in joules) released by the earthquake in this neighborhood.

SOLUTION:

